

The $d = 6$, $\mathcal{N} = 1$ heterotic string does not “live” in six dimensions[†]**Andrea Gregori¹***Humboldt-Universität, Institut für Physik
D-10115 Berlin, Germany***Abstract**

We discuss why the $\mathcal{N}_6 = 1$ heterotic string has to be viewed as something similar to a “non-compact orbifold”. Only the perturbative spectrum is forced to satisfy the constraints imposed by the vanishing of six-dimensional anomalies. These do not apply to the states of the non-perturbative spectrum, such as those appearing when small instantons shrink to zero size.

[†]Research supported by the “Marie Curie” fellowship HPMF-CT-1999-00396. Extended version of a work presented at “*Modern Trends in String Theory*”, Lisbon, 13–17 July 2001.

¹e-mail: agregori@physik.hu-berlin.de

The massless spectrum of the heterotic string in ten dimensions is determined by the constraint imposed by the vanishing of the Green–Schwarz anomaly, that requires the presence, together with the $\mathcal{N}_{10} = 1$ supergravity multiplet, of 496 gauge bosons; the only two solutions for the gauge group G are $G = E_8 \times E_8$ and $G = SO(32)$. The (massless) spectrum of the string in lower dimensions is obtained by “dimensional reduction”. Namely, the degrees of freedom remain the same, but they are differently interpreted, in terms of lower dimensional fields, arranged into multiplets of the appropriate $\mathcal{N}_d$ supersymmetry in d dimensions.

In less than ten dimensions, the Green-Schwarz anomaly constraint doesn’t apply anymore, and the gauge and matter spectrum can be varied, by introducing Wilson lines. In six dimensions, it is also possible to reduce supersymmetry from the $\mathcal{N}_6 = 2$, as derived by toroidal compactification of the $\mathcal{N}_{10} = 1$, to $\mathcal{N}_6 = 1$, by compactifying the four coordinates on a curved space. As is known, this space is unique, the K3 surface. Although not uniquely determined, the massless spectrum is nevertheless highly constrained: it must satisfy a constraint derived by requiring the cancellation of the $\mathcal{N}_6 = 1$, $\text{Tr } R^4$ anomaly. This constraint reads [1]:

$$N_H - N_V + 29N_T = 273. \quad (0.1)$$

The $\mathcal{N}_6 = 2$ heterotic string has been conjectured to be dual to the type IIA string compactified on the K3 surface [2]. When both the theories are further compactified on a two-torus, T^2 , this translates into a duality between $\mathcal{N}_4 = 4$ theories, that maps a modulus of the gravity multiplet into a modulus of the vector manifold [3]. The $\mathcal{N}_6 = 1$ theory on the other hand does not possess a type IIA dual. However, when this theory is toroidally compactified on a two-torus, it is expected to be dual to the type IIA string compactified on a K3 fibration. The heterotic dilaton–axion field maps then to the volume form of the fibration. It may seem a bit strange that, while the $\mathcal{N}_4 = 4$ duality exists also in six dimensions, this duality of the $\mathcal{N}_4 = 2$ theory appears only in four dimensions. From a technical point of view, this is related to the fact that it is not possible to construct an $\mathcal{N}_6 = 1$ type II theory. But it also means that, if string-string duality has to be taken seriously, the operation of compactification of the $\mathcal{N}_6 = 1$ heterotic theory on T^2 is not so an innocuous one: something very special must happen at the level of the “string theory”, namely the theory conjectured to be the basic one, underlying all the specific manifestations, whether heterotic or type II, or type I string constructions. From that point of view, the two-torus must not be a “flat” space.

In order to understand what is going on, we start by considering in detail the heterotic/type IIA duality map in the $\mathcal{N}_4 = 2$ theory. The correspondence of the volume form of the base of the type IIA, K3 fibration, with the heterotic dilaton can be observed by looking at the effective theory, built on the massless states of both the constructions. The duality map is therefore “perturbative/non-perturbative”, and requires for consistency that also the modulus parameterizing the coupling of the type IIA string, belonging to the hypermultiplets, has a heterotic counterpart. On the heterotic side, the hypermultiplets correspond to

moduli in the K3: the latter must therefore be elliptically fibered, in order to “contain” a torus corresponding to the type II coupling, whose dual must be a perturbative modulus of the heterotic string. For what matters instead the K3 fibration on the type IIA side, this also must satisfy certain requirements in order to constitute a compactification space compatible, in the limit of large volume of the base, with the heterotic weak coupling limit. Here however we must clarify certain commonly accepted points that we find misleading. It has often been said that the type IIA space must also be elliptically fibered. This property is advocated in order to respect a “fiberwise” identification of the type IIA and heterotic theories¹. Namely, one should read the image of the heterotic two-torus in a torus, appearing, on the type IIA side, as the base of a fibration. As we will see, the heterotic two-torus cannot be mapped into a “torus” on the type IIA side. It has also been claimed that the fiber itself should be elliptically fibered: the fiber is in fact a K3, and it should possess geometric properties corresponding to those of the K3 on the heterotic side. This also is not correct: the type IIA compactification space cannot “contain” a torus corresponding to the torus contained in the heterotic elliptic K3; this is in fact dual to the type II coupling, and the type II string cannot contain as geometric, perturbative modulus, its own coupling. In order to see what should instead be the properties of the type IIA compact space, we consider a well known example of dual pair, that of Ref. [6]. In this case, it has been possible to investigate both the string theories through the explicit construction of their partition function [7], something that allowed to perform direct computations of terms of the effective action. The image of the heterotic two-torus can therefore be followed and checked explicitly.

On the type IIA side, the model is obtained by compactification on a $CY^{11,11}$ manifold. For details about the type IIA and heterotic construction, we refer the reader to Refs. [6, 7]. Owing to the fact that $CY^{11,11}$ is self-mirror, both the vector and hyper-multiplets moduli spaces are expected to be exact, and to not receive neither perturbative nor non-perturbative corrections. In particular, $\mathcal{N}_4 = 2$ supersymmetry is expected to remain unbroken even non-perturbatively, and we can easily follow the map of the moduli of the heterotic two-torus and of the dilaton by comparing, as in Ref. [7], the corrections to the effective coupling of a term of the effective action, the R^2 term. The reason of this choice is that this quantity receives contributions only from BPS saturated multiplets, and, owing to unbroken supersymmetry, the moduli of the heterotic K3 don’t enter in the game [8, 9]. On the heterotic side we can therefore directly compute this quantity at the $T^2 \times T^4/Z_2$ orbifold limit. At this point, the Z_2 acts also as a shift in the two-torus. Moreover, this orbifold limit turns out to be appropriate because we are only interested in the map of the moduli of the torus and of the dilaton, and the Z_2 orbifold point all the Wilson lines are frozen to discrete values. As discussed in Ref. [7], this point in the heterotic moduli space corresponds to the $Z_2 \times Z_2$ orbifold point on the type IIA side. The heterotic moduli T and U , associated respectively to the Kähler class and the complex structure of the two-torus, are mapped into the moduli $T^{(i)}$, $i = 1, 2$, associated to the Kähler classes of two of the three tori left fixed by the elements of the $Z_2 \times Z_2$ orbifold group. The Kähler class of the third torus, $T^{(3)}$, is dual to the heterotic dilaton: this plane can in fact be seen as the base of a K3 fibration, whose fiber corresponds to the other two planes. The model is not symmetric in the three planes:

¹See for instance Refs. [4, 5].

the first two are acted on by a translation, dual to the translation acting on the heterotic two-torus (notice that, owing to the $Z_2 \times Z_2$ action, this space cannot be put in the trivial form of base $\times$ fiber). We learn therefore that, in this particular case:

- i) the type IIA space respects the $T \leftrightarrow U$ symmetry of the heterotic two-torus, appearing here as a symmetry between two fixed planes, $T^{(1)} \leftrightarrow T^{(2)}$, consequence of the symmetry of the action of two Z_2 orbifold elements;
- ii) the image of the three heterotic fields S, T, U , “covers” the entire type IIA compact space, namely: $\langle S \rangle \times \langle T \rangle \times \langle U \rangle \rightarrow \prod_{i=1}^3 \langle T^{(i)} \rangle \approx \text{Vol}(\text{CY})$. This is quite reasonable; the three moduli S, T, U play in fact a special role: they must map into the three two-cycles always present in the type IIA Calabi–Yau space (see Ref. [10]).

We consider now the decompactification of the heterotic torus. Here we encounter a first mismatch, in the number of coordinates that are decompactified on the two sides. Under the heterotic/type IIA map, the heterotic torus is “de-constructed” and “reconstructed” back as a four-submanifold. This “doubling” of coordinates is consistent, because on the heterotic side both the Kähler class and the complex structure of the torus belong to the moduli in the vector multiplets, while on the type II side complex structure and Kähler class deformations belong to different moduli manifolds. Therefore, part of the moduli of the type IIA four-submanifold will correspond to moduli in the heterotic K3. Moreover, although the orbifold action is free on both sides, on the heterotic side the action is trivialized by decompactifying one coordinate of the torus, while on the type IIA side this trivialization is achieved only after decompactification of two coordinates. This also is consistent, because on the heterotic side the translation in the two torus acts on both the moduli T and U , and the same must be true for its dual image. This however implies that, while on the heterotic side the orbifold action is trivialized in five dimensions, on the type IIA side it is in six dimensions. If we now decompactify the heterotic torus, we see that the dual type II picture is ill defined: if we reach the six dimensional limit by passing through five dimensions, we take the $T \rightarrow \infty$ limit while decompactifying also the field U . If instead we decompactified both the coordinates at the same time, we go to six dimensions keeping U fixed, or finite. The decompactification of the heterotic torus may therefore correspond on the type II side to a decompactification from one up to four coordinates: all these situations correspond, on the heterotic side, to the same limit ²!

From the point of view of the effective theory, the “decompactification” is quite delicate. In the case of “freely acting orbifolds”, as is the present case, there are two decompactification limits: a true, genuine decompactification, under which the orbifold projection is trivialized, and a T-dual limit, in which the orbifold operation is not trivialized [11]. In this second limit we obtain a “non-compact orbifold”, namely a space in which certain compact coordinates are “expanded”, however, from a global point of view, the space is not a plane but still an orbifold, with fixed points at which the “curvature is concentrated”: we don’t have therefore a genuine six dimensional theory, and in fact the massless degrees of

²Moreover, we saw that certain moduli in the four-submanifold of the type IIA side, that corresponds to the heterotic torus, are dual to moduli in the heterotic K3. Through heterotic/type II/heterotic duality, we see therefore that the decompactification of the heterotic torus may, by consistency, involve also a decompactification of the heterotic K3.

freedom, once interpreted in terms of states of a six dimensional theory, do not satisfy the anomaly constraint (0.1). The difference between the two situations is explicit from the very construction of the heterotic orbifold, and there is no ambiguity in understanding whether we are performing a genuine decompactification, as in the first case, or we are going to the “non-compact orbifold limit”. The translation in the torus, associated to the orbifold projection, breaks in fact T-duality explicitly and makes the two limits quite different. From the point of view of the underlying theory, in the genuine decompactification limit it is not really essential the precise identification of the space we end up on the type IIA side. We know in fact that, under this process, we obtain an approximate restoration of the initial amount of supersymmetry; in the case at hand, we get an $\mathcal{N}_4 = 4$ theory, in an appropriate decompactification limit. Since the massless spectrum of this theory was just a projection of the perturbative spectrum of the heterotic string, as derived from compactification of the ten dimensional theory, for what matters the identification of the underlying theory we deal with the process of compactification/decompactification as in a supergravity theory, with projections “à la Scherk–Schwarz”.

More subtle is the case of a true “stringy” situation. Let’s consider a heterotic theory obtained by compactification on $T^2 \times K3$, with a certain choice of gauge bundle, but without any explicit action on T^2 . Namely, a four dimensional theory obtained just by simple dimensional reduction on T^2 of a six dimensional heterotic vacuum. In general, this theory possesses a dual realization as a type IIA string compactified on an appropriate K3 fibration $\mathcal{M}$. As before, also in this case the image of the heterotic two-torus must “cover” the fiber, in order to represent, together with the dilaton field, the three “minimal” cycles of the Calabi–Yau $\mathcal{M}$, corresponding to the three “minimal” vector fields of this theory. As before, the decompactification of the heterotic torus doesn’t lead to a smooth space. However, in this case we cannot use arguments similar to those of freely acting orbifolds, for which we know that the orbifold projection is trivialized in the decompactification limit. Here, it is not anymore immaterial what is the space we end up on the type IIA side, because the final theory has the same amount of supersymmetry as the four dimensional orbifold. We are therefore in a situation quite similar to the “non compact orbifold” limit of the previous example. As long as we only consider the perturbative spectrum of the heterotic string, the compactification on a torus can be handled as in field theory: the perturbative spectrum is in fact just the “dimensional reduction” of the one obtained in six dimensions. Things are however quite different when, at special points in the hypermultiplets moduli, new massless states appear, associated to instantons that shrink to zero size [12]. These states are entirely non-perturbative from the heterotic point of view. Their existence can however be detected on the type IIA dual, where they are associated to singular cycles appearing at points in which the fiber degenerates [10]. These states do not disappear when we decompactify the image of the heterotic torus, and the “compactification” space we end up is not quite a four dimensional manifold, but a singular six dimensional space with some “directions” expanded. As it happens for “non compact orbifolds”, there is no reason to expect that, once re-interpreted in terms of six dimensions, the massless states of this theory satisfy the anomaly constraint (0.1), and in fact in general they don’t. In order to see this,

we must consider more in detail what is the structure of the type IIA Calabi–Yau space $\mathcal{M}$, dual to the heterotic string on $T^2 \times K3$. This space must possess a symmetry reflecting the $T \leftrightarrow U$ symmetry of the heterotic torus. Since, on the heterotic side, this symmetry is valid perturbatively, and can only be spoiled by instanton corrections of the type $\sim e^{-S}$, it must result on the type IIA side in a *geometric* property of the fiber: it can be spoiled only at the level of the quantum theory, by exponential terms of the type e^{-nt} , where t is the modulus parameterizing the base of the fibration. These terms are not contained in the “geometric” informations of the Calabi–Yau: they are proper of the quantum theory, and can be computed using string–string dualities, such as mirror symmetry [13, 14]. The $T \leftrightarrow U$ symmetry is therefore a symmetry of the generic fiber. For what we saw, the image of the heterotic torus is not a torus of the type IIA side, but rather something in the fiber of dimension four; this implies that the $T \leftrightarrow U$ symmetry is a “mirror” symmetry that relates a part of the fiber to another part. When, by moving in the moduli space, the fiber encounters a singularity, such that new cycles E appear, intersecting the cycle “ T ”, there exists therefore also a “mirror” singularity with cycles E' intersecting “ U ”. By this we mean that for any new triple intersection term of the type $E \cap E \cap T$, there is also a triple intersection $E' \cap E' \cap U$. From the heterotic point of view, intersections of this kind tell us that new vector multiplets appeared, with an effective coupling parameterized respectively by the volume of the two-torus, or its complex structure. Under decompactification to six dimensions, all the effective couplings are rescaled by dividing them by the volume of the two torus, $V_{(2)} \propto T_2$. This means that terms of the first kind decompactify to six dimensions to terms with coupling 1, while terms of the second kind disappear from the effective action. Being however the two situations related by T-duality in one coordinate, it is easy to realize that the second type of vectors decompactify to tensors in six dimensions. We have therefore the following situation: once decompactified to six dimensions, the perturbative spectrum decompactifies as usual to a spectrum that satisfies the constraint (0.1). The extra states give rise to an equal number of extra vector and tensor multiplets. There is no choice of extra hypermultiplets for which the additional states can satisfy the constraint (0.1): their contribution appears in fact with the same sign as the tensor multiplets.

Terms with six dimensional coupling 1 have been investigated in several works [15, 16, 17]. These analysis of non-perturbative states of the heterotic string are however incomplete: they neglect the fact that the $\mathcal{N}_6 = 1$ theory is not a “truly six-dimensional” theory. In taking flat limits ($K3 \rightarrow R^4$) on which to study small instantons configurations, one always implicitly chooses a preferred direction of decompactification, treating the image of the heterotic two-torus on the type IIA side as a sub-manifold of dimension two instead of four. Moreover, any constraint imposed by the vanishing of six dimensional anomalies cannot be applied, being completely meaningless because we are not really in six dimensions. In particular, it is not necessary to introduce couplings of the extra tensors with Chern–Simons terms, whose existence would be required by six dimensional anomaly cancellation. Analogously, incomplete are also the informations about the heterotic non-perturbative spectrum obtained by looking at the perturbative spectrum of the type I string with D9 and D5 branes.

In our discussion we made use of string–string duality in order to make clear that the $\mathcal{N}_6 = 1$ heterotic string doesn’t really live in six dimensions. However, we don’t really need

heterotic/type II duality in order to see that the non-perturbative spectrum violates the constraint (0.1). This can be seen already from the heterotic string itself. Let's in fact suppose that, at certain points in the K3 moduli space new massless states appear, with coupling 1 in six dimensions. Under compactification to four dimensions, their effective coupling becomes then $1/g_{(4)}^2 \propto V_{(\text{torus})} \equiv \text{Im } T$. The absence of “mirror” states with coupling $\text{Im } U$ explicitly contradicts the $T \leftrightarrow U$ symmetry, direct consequence of T-duality and toroidal compactification. The six dimensional theory must therefore contain also these states, that can only be tensors; (0.1) is then automatically violated. We could on the other hand suppose that T-duality is preserved because, under compactification, in some misterious way the coupling “1” of six dimensions acquires, besides the volume factor $\text{Im } T$, also a dependence on U : $1/g_{(4)}^2 \approx \text{Im } T + \text{Im } U$ (to be eventually promoted to an $SL(2, Z)$ -invariant function of these fields). Such an expression is compatible with the coupling of six dimensions, because the U -dependent part vanishes anyway under decompactification and rescaling with the volume of the two-torus. However, with such a coupling, from the four dimensional point of view we could not discriminate whether the new vector fields originate from six dimensional vectors or tensors: we would have an effective doubling of kinetic and interaction terms, and therefore get anyway into a contradiction. In summary,

perturbative T-duality and the appearance of non-perturbative massless states necessarily implies that the six dimensional anomaly constraint (0.1) is violated.

We stress that this is peculiar of string theory, and does not happen in field theory/supergravity: T-duality and non-perturbative states of the “small instantons” type are pure stringy phenomena.

As a last comment, we remark that, always by string-string duality, we learn that also the ten dimensional $\mathcal{N}_{10} = 1$ heterotic string is in some sense a “non-compact orbifold”. This can be seen also by considering it as a Hořava–Witten orbifold: when a further circle of the M-theory is compactified to a small radius, we fall into a perturbative type IIA string situation, for which it is not possible to project with a Z_2 twist only one coordinate. However, in this case there are no non-perturbative extensions of the heterotic massless spectrum; therefore, one can safely neglect such details and deal with the string massless spectra as in field theory.

References

- [1] J. H. Schwarz, *Anomaly-Free Supersymmetric Models in Six Dimensions*, Phys. Lett. **B371** (1996) 223–230, [hep-th/9512053](#).
- [2] E. Witten, *String theory dynamics in various dimensions*, Nucl. Phys. **B443** (1995) 85–126, [hep-th/9503124](#).
- [3] C. M. Hull and P. K. Townsend, *Unity of superstring dualities*, Nucl. Phys. **B438** (1995) 109–137, [hep-th/9410167](#).
- [4] P. S. Aspinwall, *K3 surfaces and string duality*, [hep-th/9611137](#).

- [5] P. S. Aspinwall and M. R. Plesser, *T-duality can fail*, JHEP **08** (1999) 001, [hep-th/9905036](#).
- [6] S. Ferrara, J. A. Harvey, A. Strominger, and C. Vafa, *Second quantized mirror symmetry*, Phys. Lett. **B361** (1995) 59–65, [hep-th/9505162](#).
- [7] A. Gregori, C. Kounnas, and P. M. Petropoulos, *Non-perturbative gravitational corrections in a class of $N = 2$ string duals*, Nucl. Phys. **B537** (1999) 317–343, [hep-th/9808024](#).
- [8] J. A. Harvey and G. Moore, *Algebras, BPS States, and Strings*, Nucl. Phys. **B463** (1996) 315–368, [hep-th/9510182](#).
- [9] J. A. Harvey and G. Moore, *Exact gravitational threshold correction in the FHSV model*, Phys. Rev. **D57** (1998) 2329–2336, [hep-th/9611176](#).
- [10] P. S. Aspinwall and J. Louis, *On the Ubiquity of $K3$ Fibrations in String Duality*, Phys. Lett. **B369** (1996) 233–242, [hep-th/9510234](#).
- [11] E. Kiritsis, C. Kounnas, P. M. Petropoulos, and J. Rizos, *Solving the decompactification problem in string theory*, Phys. Lett. **B385** (1996) 87–95, [hep-th/9606087](#).
- [12] E. Witten, *Small Instantons in String Theory*, Nucl. Phys. **B460** (1996) 541–559, [hep-th/9511030](#).
- [13] P. Candelas, X. C. D. L. Ossa, P. S. Green, and L. Parkes, *A pair of Calabi-Yau manifolds as an exactly soluble superconformal theory*, Nucl. Phys. **B359** (1991) 21–74.
- [14] P. Candelas, X. C. D. la Ossa, P. S. Green, and L. Parkes, *An Exactly soluble superconformal theory from a mirror pair of Calabi-Yau manifolds*, Phys. Lett. **B258** (1991) 118–126.
- [15] D. R. Morrison and C. Vafa, *Compactifications of F-Theory on Calabi-Yau Threefolds – I*, Nucl. Phys. **B473** (1996) 74–92, [hep-th/9602114](#).
- [16] D. R. Morrison and C. Vafa, *Compactifications of F-Theory on Calabi-Yau Threefolds – II*, Nucl. Phys. **B476** (1996) 437–469, [hep-th/9603161](#).
- [17] P. S. Aspinwall and D. R. Morrison, *Point-like instantons on $K3$ orbifolds*, Nucl. Phys. **B503** (1997) 533–564, [hep-th/9705104](#).